

Normality via local spectral radii

Dariusz Cichoń, Il Bong Jung, and Jan Stochel

IMUJ PREPRINT 2007/08

ABSTRACT. A new criterion for essential normality of unbounded Hilbert space operators is furnished in terms of local spectral radius. Accordingly, extensive study of operators of certain types related to local spectral radius is conducted. Spectral radii of local restrictions of a normal operator are investigated.

1. Introduction. In this paper, we give conditions for normality of the closure (read: essential normality) of an unbounded Hilbert space operator. Our approach is based on the notion of local spectral radius (cf. [7, 35, 36, 8, 19, 37]; see also [4, 5] for the case of unbounded operators). We attach to any (unbounded) operator a family of invariant linear spaces of C^∞ -vectors determined by uniform boundedness of their local spectral radii. Examining the corresponding restrictions (called local restrictions) of the operator in question enables us to distinguish classes of localoid and locally normaloid operators (cf. Section 5). This is preceded by detailed analysis of local spectral radius, as some of its properties are not preserved in the unbounded case (cf. Sections 2 and 4). In Section 3, we provide necessary and sufficient conditions for an abstract transformation of the half real line to come from spectral radii of local restrictions of a normal operator. The advantage of introducing notions of localoid and locally normaloid operators when compared with other known classes of operators (e.g. restriction-normaloid operators) lies in the fact that

1991 *Mathematics Subject Classification.* Primary 47B15, 47B20; Secondary 47B07.

Key words and phrases. local spectral radius, normal operator, paranormal operator, normaloid operator, algebraic operator, locally algebraic operator, compact operator.

The first and the third authors were supported by the KBN grant 2 P03A 037 024. The second author was supported by a grant (R14-2003-006-01000-0) from the Korea Research Foundation.

our attention is confined to the family of selected invariant subspaces which may reduce to a finite set. We show that paranormal operators as well as those satisfying the Kato-Protter inequality form proper subclasses of locally normaloid operators. In Section 6 we discuss the question of when essential normality of an operator follows from essential normality of all its cyclic parts. Having this established, we provide a criterion for essential normality of locally algebraic operators written in terms of local normaloidity (cf. Theorem 34).

2. Local spectral radius. Throughout what follows, \mathcal{H} stands for a complex Hilbert space. By an operator *in* \mathcal{H} we mean a linear mapping $A: \mathcal{D}(A) \rightarrow \mathcal{H}$ defined on a linear subspace $\mathcal{D}(A)$ of \mathcal{H} , called the *domain* of A . We denote by $\mathcal{N}(A)$, A^* and \bar{A} the kernel, the adjoint and the closure of A respectively. We put $\mathcal{D}^\infty(A) = \bigcap_{n=1}^\infty \mathcal{D}(A^n)$. A densely defined operator A in \mathcal{H} is called *normal* if it is closed and $A^*A = AA^*$. An operator A in \mathcal{H} is said to be *essentially normal* if A is closable and \bar{A} is normal.

Given an inner product space \mathcal{D} , we denote by $\mathbf{B}(\mathcal{D})$ the algebra of all bounded linear mappings from \mathcal{D} into \mathcal{D} . If $A \in \mathbf{B}(\mathcal{D})$, then $r(A)$ stands for the spectral radius of the unique bounded linear extension of A to the completion of \mathcal{D} ; in the case of $\mathcal{D} = \{0\}$, we put $r(A) = 0$. By the Gelfand theorem $r(A) = \lim_{n \rightarrow \infty} \|A^n\|^{1/n}$.

Given an operator A in \mathcal{H} , we define

$$r(A, f) = \limsup_{n \rightarrow \infty} \|A^n f\|^{1/n}, \quad f \in \mathcal{D}^\infty(A),$$

$$\mathcal{H}_A(t) = \{f \in \mathcal{D}^\infty(A) : r(A, f) \leq t\}, \quad t \in [0, \infty].$$

The quantity $r(A, f) \in [0, \infty]$ is called the *local spectral radius* of A at f . Note that the set of all vectors $f \in \mathcal{H}$ such that $r(A, f) < \infty$ is identical with the set of all bounded vectors for A (cf. [10, 32] and [33] for an up-to-date approach). Evidently $\mathcal{H}_A(s) \subseteq \mathcal{H}_A(t)$ whenever $0 \leq s \leq t$, and $\mathcal{H}_A(\infty) = \mathcal{D}^\infty(A)$. Let us list some basic properties of local spectral radius (consult the proof of [8, Lemma 1]).

LEMMA 1. *If A is an operator in \mathcal{H} , then*

- (i) $r(A, \alpha f + \beta g) \leq \max\{r(A, f), r(A, g)\}$ for all $\alpha, \beta \in \mathbb{C}$ and $f, g \in \mathcal{D}^\infty(A)$,
- (ii) $r(A, Af) = r(A, f)$ for all $f \in \mathcal{D}^\infty(A)$,
- (iii) for every $t \in [0, \infty]$, $\mathcal{H}_A(t)$ is a linear subspace of $\mathcal{D}^\infty(A)$ which is invariant for A ,
- (iv) for every $t \in [0, \infty]$, $\mathcal{H}_A(t) = \mathcal{H}_A(\tilde{t})$, where $\tilde{t} \stackrel{\text{def}}{=} \sup\{r(A, f) : f \in \mathcal{H}_A(t)\}$.

For $t \in [0, \infty]$, we put $A_{[t]} = A|_{\mathcal{H}_A(t)}$. We regard $A_{[t]}$ as an operator in \mathcal{H} .

We now show to what extent the equality $r(A, f)^k = r(A^k, f)$, originally proved in [8], can be extended to the case of unbounded operators.

LEMMA 2. *If A is an operator in \mathcal{H} , f is a vector in $\mathcal{D}^\infty(A)$ and $k \geq 1$ is an integer, then*

- (i) $r(A, f)^k = \max\{r(A^k, A^j f) : j = 0, 1, \dots, k-1\}$,
- (ii) $r(A, f)^k = \max\{r(A^k, A^j f) : j = 0, 1\}$ provided $A_{[t]}$ is closed (as an operator in \mathcal{H}) for $t = r(A, f)$,
- (iii) $r(A, f)^k = r(A^k, f)$ provided $A_{[t]}$ is bounded for $t = r(A, f)$.

PROOF. (i) Each positive integer n can be written as $n = km + j$ with unique nonnegative integers $m \geq 0$ and $j \in \{0, \dots, k-1\}$. This leads to

$$\begin{aligned} r(A, f)^k &= \lim_{l \rightarrow \infty} \sup_{n \geq l} \|A^n f\|^{k/n} \\ &= \lim_{l \rightarrow \infty} \max_{j=0, \dots, k-1} \sup_{km+j \geq l} \left(\|(A^k)^m A^j f\|^{1/m} \right)^{1/(1+\frac{j}{km})} \\ &= \max_{j=0, \dots, k-1} \lim_{l \rightarrow \infty} \sup_{km+j \geq l} \left(\|(A^k)^m A^j f\|^{1/m} \right)^{1/(1+\frac{j}{km})} \\ &= \max_{j=0, \dots, k-1} r(A^k, A^j f). \end{aligned}$$

(ii) Set $\|h\|_A = \sqrt{\|h\|^2 + \|Ah\|^2}$ for $h \in \mathcal{H}_A(t)$. Denote by \mathcal{X}_t the inner product space $(\mathcal{H}_A(t), \|\cdot\|_A)$ and define the linear mapping $B_t: \mathcal{X}_t \rightarrow \mathcal{X}_t$ by $B_t h = Ah$ for $h \in \mathcal{X}_t$. Since $A_{[t]}$ is a closed operator in \mathcal{H} , \mathcal{X}_t is a Hilbert space and the operator B_t is closed. By the closed graph theorem $B_t \in \mathbf{B}(\mathcal{X}_t)$. Note that $r(A, f) = r(B_t, f)$ and $r(B_t^k, f) = \max\{r(A^k, f), r(A^k, Af)\}$ ($r(B_t, f)$ and $r(B_t^k, f)$ are calculated with respect to $\|\cdot\|_A$). Applying [8, Lemma 1 (6)] to the bounded operator B_t we get

$$r(A, f)^k = r(B_t, f)^k = r(B_t^k, f) = \max\{r(A^k, f), r(A^k, Af)\}.$$

(iii) Applying [8, Lemma 1 (6)] now to the bounded operator $A_{[t]}$, we get

$$r(A, f)^k = r(A_{[t]}, f)^k = r(A_{[t]}^k, f) = r(A^k, f), \quad k \geq 1,$$

which completes the proof. \square

Note that the case $r(A, f) = \infty$ is excluded in (ii) and (iii) of Lemma 2.

COROLLARY 3. *If A is an operator in \mathcal{H} such that all the operators $A_{[t]}$, $t \in [0, \infty)$, are bounded, then $\mathcal{H}_{A^k}(t) = \mathcal{H}_A(\sqrt[k]{t})$ for all real $t \geq 0$ and all integers $k \geq 1$.*

COROLLARY 4. *If A is an operator in \mathcal{H} and $k \geq 1$ is an integer, then*

(i) *for every $f \in \mathcal{D}^\infty(A)$ there exists $j \in \{0, \dots, k-1\}$ such that*

$$r(A, A^j f)^k = r(A^k, A^j f),$$

(ii) *if $\mathcal{D}^\infty(A) \neq \{0\}$, then there exists $f \in \mathcal{D}^\infty(A) \setminus \{0\}$ such that $r(A, f)^k = r(A^k, f)$.*

PROOF. (i) follows from Lemmata 1 and 2.

(ii) Take any nonzero vector $f \in \mathcal{D}^\infty(A)$. If $A^k f = 0$, then manifestly $r(A, f)^k = r(A^k, f) = 0$. Otherwise, $A^j f \neq 0$ for all $j = 0, \dots, k-1$, which combined with (i) completes the proof. \square

EXAMPLE 5. We show that the equality $r(A, f)^k = r(A^k, f)$ is no longer true for arbitrary unbounded operators. Let $\{e_n\}_{n=0}^\infty$ be an orthonormal basis of \mathcal{H} and \mathcal{D} be the linear span of $\{e_n\}_{n=0}^\infty$. Given a sequence $\{\lambda_n\}_{n=0}^\infty \subseteq (0, \infty)$, we define the operator A in \mathcal{H} by $\mathcal{D}(A) = \mathcal{D}$ and $Ae_n = \lambda_n e_{n+1}$ for all integers $n \geq 0$.

Consider the sequence $(\lambda_0, \lambda_1, \lambda_2, \dots) = (u, u^{-1}, u^2, u^{-2}, u^3, u^{-3}, \dots)$, where $u > 1$ is a fixed real number. It follows from Lemma 1 that

$$(1) \quad r(A, e_j) = r(A, A^j e_0) = r(A, e_0) = \sqrt{u}, \quad j \geq 1.$$

Take $f = \sum_{l=0}^N \alpha_l e_l \in \mathcal{D} \setminus \{0\}$ with $\{\alpha_l\}_{l=0}^N \subseteq \mathbb{C}$. By Lemma 1 (i) we have

$$(2) \quad r(A, f) \leq \max\{r(A, e_j) : j = 0, \dots, N\} \stackrel{(1)}{=} \sqrt{u},$$

$$(2) \quad r(A, f) = \limsup_{n \rightarrow \infty} \left(\sum_{l=0}^N \|\alpha_l A^n e_l\|^2 \right)^{1/(2n)} \geq \limsup_{n \rightarrow \infty} \|\alpha_p A^n e_p\|^{1/n} = \sqrt{u},$$

where p is chosen so that $\alpha_p \neq 0$. As a consequence, we obtain

$$(3) \quad r(A, f) = \sqrt{u}, \quad f \in \mathcal{D} \setminus \{0\}.$$

Arguing as above we see that

$$(4) \quad r(A^2, f) = \begin{cases} u & \text{if there exists an odd integer } p \text{ such that } \alpha_p \neq 0, \\ 1 & \text{otherwise.} \end{cases}$$

The equalities (3) and (4) imply that

$$(5) \quad \mathcal{H}_A(t) = \begin{cases} \{0\} & \text{if } t < \sqrt{u}, \\ \mathcal{D} & \text{if } t \geq \sqrt{u}, \end{cases} \quad \text{and} \quad \mathcal{H}_{A^2}(t) = \begin{cases} \{0\} & \text{if } t < 1, \\ * & \text{if } 1 \leq t < u, \\ \mathcal{D} & \text{if } t \geq u, \end{cases}$$

where “*” stands for the linear span of $\{e_{2j}\}_{j=0}^{\infty}$.

In turn, if $(\lambda_0, \lambda_1, \lambda_2, \dots) = (2^{2^0}, 2^{-2^0}, 2^{2^1}, 2^{-2^1}, 2^{2^2}, 2^{-2^2}, \dots)$, then one can show that equalities (3), (4) and (5) remain valid with ∞ in place of u and \sqrt{u} .

We now list some basic properties of the function $t \mapsto r(A_{[t]})$.

LEMMA 6. *If A is an operator in \mathcal{H} such that $A_{[u]} \in \mathbf{B}(\mathcal{H}_A(u))$ for some $u \in [0, \infty)$, then*

- (i) $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ for all $t \in [0, u]$,
- (ii) $\mathcal{H}_A(t) \subseteq \mathcal{H}_A(r(A_{[t]}))$ for all $t \in [0, u]$,
- (iii) $r(A_{[t]}) \leq r(A_{[t']})$ whenever $t \in [0, u]$ and $A_{[t']} \in \mathbf{B}(\mathcal{H}_A(t'))$, where $t' \stackrel{\text{def}}{=} r(A_{[t]})$,
- (iv) $t \leq r(A_{[t]})$ whenever $t \in [0, u]$ and $t = r(A, f)$ for some $f \in \mathcal{D}^\infty(A)$,
- (v) the function $[0, u] \ni t \mapsto r(A_{[t]}) \in [0, \infty)$ is monotonically increasing,
- (vi) if $t \in [0, u)$ and $r(A_{[t]}) < r(A_{[t+]}) \stackrel{\text{def}}{=} \lim_{s \rightarrow t^+} r(A_{[s]})$, then $t < r(A_{[s]})$ for all $s \in (t, u]$ and consequently $t \leq r(A_{[t+]})$,
- (vii) if $t \in (0, u]$ and $r(A_{[t]}) > r(A_{[t-]}) \stackrel{\text{def}}{=} \lim_{s \rightarrow t^-} r(A_{[s]})$, then $r(A_{[t]}) \geq t$.

PROOF. (i) Evident.

(ii) This follows from the inequality

$$(6) \quad r(A, f) = r(A_{[t]}, f) \leq r(A_{[t]}), \quad f \in \mathcal{H}_A(t), \quad t \in [0, u].$$

(iii)&(v) Apply (ii) and the Gelfand formula for the spectral radius.

(iv) This is a direct consequence of (6).

(vi) Suppose, for contradiction, that (vi) fails to hold. Then there exist $t \in [0, u)$ and $s \in (t, u]$ such that $r(A_{[t]}) < r(A_{[t+]})$ and $r(A_{[s]}) \leq t$. By (ii) this yields

$$(7) \quad \mathcal{H}_A(s) \subseteq \mathcal{H}_A(r(A_{[s]})) \subseteq \mathcal{H}_A(t) \subseteq \mathcal{H}_A(s).$$

Hence $r(A_{[t]}) = r(A_{[s]})$ and thus by (v), $r(A_{[t]}) = r(A_{[t+]})$, which is a contradiction.

(vii) Since $r(A_{[t]}) > r(A_{[t-]})$, we see that $\mathcal{H}_A(s) \subsetneq \mathcal{H}_A(t)$ for every $s \in (0, t)$. Hence for every $s \in (0, t)$ there exists $f_s \in \mathcal{D}^\infty(A)$ such that $s < r(A, f_s) \leq t$. Observing that $r(A, f_s) \leq r(A_{[t]})$ and letting s tend to t , we get $t \leq r(A_{[t]})$. \square

3. Characterization of $t \mapsto r(A_{[t]})$ for normal operators. The function $t \mapsto r(A_{[t]})$ can be explicitly computed for normal operators by means of their spectral measures. For $t \in [0, \infty)$, we put $\Delta_t = \{z \in \mathbb{C}: |z| \leq t\}$, Δ_t° = the interior of Δ_t and $\Gamma_t = \Delta_t \setminus \Delta_t^\circ$. By convention $\sup \emptyset = 0$ and $|Y| = \{|z|: z \in Y\}$ for $Y \subseteq \mathbb{C}$. As usual, $\sigma(A)$ stands for the spectrum of an operator A .

LEMMA 7. *If E is the spectral measure of a normal operator A in \mathcal{H} , then for all $t \in [0, \infty)$,*

- (i) $\mathcal{H}_A(t) = E(\Delta_t)\mathcal{H}$,
- (ii) $r(A_{[t]}) = \begin{cases} t & \text{if } E(\Gamma_t) \neq 0, \\ \sup |\sigma(A) \cap \Delta_t^\circ| & \text{if } E(\Gamma_t) = 0, \end{cases}$
- (iii) $r(A_{[t]})$ equals $\sup |\sigma(A) \cap \Delta_t^\circ|$ or $\sup |\sigma(A) \cap \Delta_t|$.

PROOF. (i) Consult the proof of [33, Proposition 4].

(ii) & (iii) Take $t \in [0, \infty)$. By (i), $E_{[t]}(\cdot) \stackrel{\text{def}}{=} E(\cdot)|_{E(\Delta_t)\mathcal{H}}$ is the spectral measure of $A_{[t]}$. Since the closed support of the spectral measure of a normal operator coincides with its spectrum, we get

$$(8) \quad \sigma(A_{[t]}) \subseteq \Delta_t \cap \sigma(A) \quad \text{and} \quad \sigma(A_{[t]}) \cap \Delta_t^\circ = \sigma(A) \cap \Delta_t^\circ.$$

Suppose that $E(\Gamma_t) \neq 0$. The standard compactness argument implies that there exists $z_0 \in \Gamma_t$ such that $E(U \cap \Gamma_t) \neq 0$ and consequently $E(U \cap \Delta_t) \neq 0$ for every open neighborhood U of z_0 . This means that z_0 belongs to the closed support of $E_{[t]}$. Hence, by (8), $r(A_{[t]}) = t$. As a consequence, $r(A_{[t]}) = \sup |\sigma(A) \cap \Delta_t|$.

Suppose now that $E(\Gamma_t) = 0$. If $\sup |\sigma(A) \cap \Delta_t^\circ| = t$, then by (8), $r(A_{[t]}) = t$ and consequently $r(A_{[t]}) = \sup |\sigma(A) \cap \Delta_t|$. In turn, if $s \stackrel{\text{def}}{=} \sup |\sigma(A) \cap \Delta_t^\circ| < t$, then $E(\Delta_t^\circ \setminus \Delta_s) = 0$. Since moreover $E(\Gamma_t) = 0$, we get $E_{[t]}(\mathbb{C} \setminus \Delta_s) = 0$, which yields $r(A_{[t]}) \leq s$. If $s = 0$, then evidently $r(A_{[t]}) = s$. Otherwise $s > 0$ and consequently $\Gamma_s \cap \sigma(A) \neq \emptyset$. This combined with (8) and $r(A_{[t]}) \leq s$ leads to $r(A_{[t]}) = s$. \square

We now extract some properties of the function $s \mapsto r(A_{[s]})$, where A is a normal operator. Their abstract versions determine all functions of this type (cf. Theorem 10). In what follows, we preserve the notation used in Lemma 6.

PROPOSITION 8. *If A is a normal operator in \mathcal{H} and $t \in [0, \infty)$, then*

- (i) $r(A_{[t]})$ equals $r(A_{[t-]})$ or $r(A_{[t+]})$,
- (ii) if the function $s \mapsto r(A_{[s]})$ is not continuous at t , then

$$\begin{aligned} r(A_{[t-]}) &< r(A_{[t+]}) = t, \\ r(A_{[v]}) &= r(A_{[t-]}), \quad v \in [r(A_{[t-]}), t), \end{aligned}$$

- (iii) if $r(A_{[s]}) = t$ for all $s \in (t, \infty)$, then $r(A_{[t]}) = t$,
- (iv) if the function $s \mapsto r(A_{[s]})$ is continuous on an open interval $U \subseteq (0, \infty)$, then there exists $c \in [0, \infty]$ such that $r(A_{[s]}) = \min\{s, c\}$ for all $s \in U$.

PROOF. It follows from Lemma 7 (ii) that

$$(9) \quad r(A_{[s]}) \leq s, \quad s \in [0, \infty).$$

The next step of the proof is to show that for every $x \in [0, \infty)$,

$$(10) \quad \text{if } r(A_{[x]}) < x, \text{ then } r(A_{[v]}) = r(A_{[x]}) \text{ for all } v \in [r(A_{[x]}), x].$$

Indeed, if $r(A_{[x]}) < x$, then by Lemma 7(ii), $u \stackrel{\text{def}}{=} \sup |\sigma(A) \cap \Delta_x^\circ| < x$ and $E(\Delta_x \setminus \Delta_u) = 0$. This and again Lemma 7(ii) give $r(A_{[v]}) = u$ for all $v \in (u, x]$. Consequently, $r(A_{[x]}) = u$. Hence it remains to verify that $r(A_{[u]}) = u$. Suppose that, contrary to our claim, $z \stackrel{\text{def}}{=} r(A_{[u]}) < u$. Then, by Lemma 7(ii), $E(\Delta_u \setminus \Delta_z) = 0$. This and $E(\Delta_x^\circ \setminus \Delta_u) = 0$ yield $E(\Delta_x^\circ \setminus \Delta_z) = 0$, which in turn implies that $\sigma(A) \cap (\Delta_x^\circ \setminus \Delta_z) = \emptyset$. Thus $\Gamma_u \cap \sigma(A) = \emptyset$, which contradicts $\sup |\sigma(A) \cap \Delta_x^\circ| = u > 0$.

We are now ready to prove all the parts of the conclusion using only (9), (10) and the monotonicity of $s \mapsto r(A_{[s]})$ (without any recourse to Lemma 7). This observation is essential in the context of the proof of Proposition 11.

(i) If $r(A_{[t]}) = t$, then by (9) and the monotonicity of the function $s \mapsto r(A_{[s]})$ we have $r(A_{[t]}) = r(A_{[t+]})$. Otherwise $r(A_{[t]}) < t$, which, together with (10), yields $r(A_{[t]}) = r(A_{[t-]})$.

(ii) By discontinuity of the function $s \mapsto r(A_{[s]})$ at t , $r(A_{[t-]}) < r(A_{[t+]}) \leq t$. Let $\{t_n\}_{n=1}^\infty \subseteq (r(A_{[t-]}), t)$ be a strictly increasing sequence tending to t . Take $v \in [r(A_{[t-]}), t)$. Since $r(A_{[t_n]}) \leq r(A_{[t-]})$ for all $n \geq 1$ and $t_n \nearrow t$, there exists an integer $k \geq 1$ such that $v \in [r(A_{[t_n]}), t_n)$ for all $n \geq k$. In virtue of (10) we see that $r(A_{[v]}) = r(A_{[t_n]})$ for all $n \geq k$. Letting n tend to ∞ , we get $r(A_{[v]}) = r(A_{[t-]})$.

We now show that $r(A_{[t+]}) = t$. In the contrary case, $r(A_{[t+]}) < t$ and consequently there exists real $s > t$ such that $r(A_{[s]}) \in [r(A_{[t+]})], t)$. This leads to $r(A_{[s]}) < t < s$. By (10), $r(A_{[v]}) = r(A_{[s]})$ for all $v \in (r(A_{[s]}), t)$. Therefore, we have $r(A_{[t-]}) = r(A_{[s]}) \geq r(A_{[t+]})$, which is a contradiction.

(iii) Apply (10) to some $x > t$.

(iv) We claim that if $r(A_{[v]}) < v$ for some $v \in U$, then $r(A_{[s]}) = r(A_{[v]})$ for all $s \in (r(A_{[v]}), \infty) \cap U$. By (10) we see that $r(A_{[s]}) = r(A_{[v]})$ for all $s \in (r(A_{[v]}), v]$. Suppose that, contrary to our claim, the function $s \mapsto r(A_{[s]})$ is not constant on $(r(A_{[v]}), \infty) \cap U$. Then there exists $w \in U$ such that $v < w$ and $r(A_{[v]}) < r(A_{[w]})$. By the Darboux property we can assume without loss of generality that $r(A_{[w]}) < v$. Hence $v \in (r(A_{[w]}), w)$, which by (10) contradicts $r(A_{[v]}) < r(A_{[w]})$.

We now show that if $r(A_{[v]}) = v$ for some $v \in U$, then $r(A_{[s]}) = s$ for all $s \in (0, v] \cap U$. Indeed, supposing that $r(A_{[w]}) \neq w$ for some $w \in (0, v) \cap U$, we infer from (9) that $r(A_{[w]}) < w$, which contradicts the previous paragraph.

To complete the proof, note that if $r(A_{[s]}) = s$ for all $s \in U$, then we may take $c = \infty$. Otherwise, there exists $v \in U$ such that $r(A_{[v]}) < v$, and so $c \stackrel{\text{def}}{=} r(A_{[v]})$ fits into (iv). This finishes the proof. \square

Given $a = \{a_n\}_{n=1}^\infty \subseteq \mathbb{C}$, we denote by M_a the *diagonal* operator in ℓ^2 defined by

$$\mathcal{D}(M_a) = \left\{ \{x_n\}_{n=1}^\infty \in \ell^2 : \sum_{n=1}^\infty |a_n x_n|^2 < \infty \right\} \quad \text{and}$$

$$M_a(x) = \{a_n x_n\}_{n=1}^\infty \quad \text{for } x = \{x_n\}_{n=1}^\infty \in \mathcal{D}(M_a).$$

It is well known that M_a is normal and $\sigma(M_a)$ is the closure of the set $\{a_n : n \geq 1\}$.

COROLLARY 9. *If $a = \{a_n\}_{n=1}^\infty \subseteq \mathbb{C}$, then*

$$(11) \quad r((M_a)_{[t]}) = \sup\{|a_n| : n \geq 1, |a_n| \leq t\}, \quad t \in [0, \infty).$$

PROOF. The spectral measure E of M_a is given by $E(\sigma) = M_{a_\sigma}$, where $a_\sigma = \{a_{\sigma,n}\}_{n=1}^\infty$ is the sequence given by $a_{\sigma,n} = 1$ if $a_n \in \sigma$ and $a_{\sigma,n} = 0$ otherwise. Thus $E(\sigma) \neq 0$ if and only if $\sigma \cap \{a_n: n \geq 1\} \neq \emptyset$. Applying Lemma 7(ii) and the equality $\sup |\sigma(A) \cap \Delta_t^o| = \sup\{|a_n|: n \geq 1, |a_n| < t\}$ completes the proof. \square

We now show that monotonically increasing transformations of $[0, \infty)$ satisfying abstract versions of conditions (i)–(iv) of Proposition 8 come from normal operators.

THEOREM 10. *If $\varphi: [0, \infty) \rightarrow [0, \infty)$ is a monotonically increasing function satisfying the following four conditions:*

- 1° *for every $t \in [0, \infty)$, $\varphi(t)$ equals $\varphi(t-)$ or $\varphi(t+)$,*
- 2° *for every $t \in (0, \infty)$, if the function φ is not continuous at t , then $\varphi(t-) < \varphi(t+) = t$ and $\varphi(v) = \varphi(t-)$ for all $v \in [\varphi(t-), t)$,*
- 3° *for every $t \in [0, \infty)$, if $\varphi(s) = t$ for all $s \in (t, \infty)$, then $\varphi(t) = t$,*
- 4° *if the function φ is continuous on an open interval $U \subseteq (0, \infty)$, then there exists $c \in [0, \infty]$ such that $\varphi(s) = \min\{s, c\}$ for all $s \in U$,*

then there exists a positive selfadjoint operator A in ℓ^2 such that $\varphi(s) = r(A_{[s]})$ for all $s \in [0, \infty)$.

PROOF. In what follows, K' stands for the collection of all accumulation points of a set $K \subseteq \mathbb{R}$. Define the following four sets:

$$\begin{aligned} K_- &= \{t \in [0, \infty): \varphi(t) = \varphi(t-) < \varphi(t+)\}, \\ K_+ &= \{t \in [0, \infty): \varphi(t-) < \varphi(t+) = \varphi(t)\}, \\ K_c &= \{t \in [0, \infty): \varphi(t-) = \varphi(t) = \varphi(t+)\}, \\ K_r &= \{t \in [0, \infty): \varphi^{-1}(\{t\}) \text{ has a nonempty interior}\}. \end{aligned}$$

It follows from 1° that $[0, \infty) = K_- \cup K_+ \cup K_c$ with pairwise disjoint terms. Clearly, $K_r \subseteq \varphi([0, \infty))$ and, by 2°, $K_+ \subseteq \varphi([0, \infty))$. It is also easily seen that the sets K_- , K_+ and K_r are at most countable. This implies that there exists a sequence $a = \{a_n\}_{n=1}^\infty \subseteq \varphi([0, \infty))$ dense in $\overline{\varphi([0, \infty))}$ such that $K_+ \cup K_r \subseteq \{a_n: n \geq 1\}$. Let A be the diagonal operator M_a . Since A is positive and selfadjoint, it remains to show that $\varphi(s) = r(A_{[s]})$ for all $s \in [0, \infty)$. We split the proof into several steps. The monotonicity assumption will be mentioned explicitly only in more subtle cases.

STEP 1. $\varphi(s) \leq s$ for all $s \in [0, \infty)$ (an analogue of (9)).

This can be done by analyzing three possible cases: $s \in K_- \cup K_+$, $s \in K_c \cap (K_- \cup K_+)'$ and $s \in K_c \setminus (K_- \cup K_+)'$.

STEP 2. If $t \in K_r$, then either $\varphi^{-1}(\{t\}) = [t, \infty)$ or there exists $u \in (t, \infty) \cap (K_- \cup K_+)$ such that $[t, u] \subseteq \varphi^{-1}(\{t\}) \subseteq [t, u]$.

Taking any nonempty open interval $(\alpha, \beta) \subseteq \varphi^{-1}(\{t\})$, we infer from Step 1 and 2° that $\alpha \in [t, \infty)$ and $(t, \beta) \subseteq K_c$, which enables us to define

$$u = \sup\{s \in (t, \infty): (t, s) \subseteq K_c\}.$$

Clearly, $u \geq \beta$ and $(t, u) \subseteq K_c$. If $u = \infty$, then 4° and 3° give $\varphi^{-1}(\{t\}) = [t, \infty)$. Assume that $u < \infty$. It follows from 4° that $(t, u) \subseteq \varphi^{-1}(\{t\})$. If $u \in K_- \cup K_+$, then applying 2° we get the other part of the conclusion of Step 2. We are left with verifying that the remaining possibility $u \in K_c$ can never happen. Indeed, the case $u \in K_c \setminus (K_- \cup K_+)'$ immediately contradicts the definition of u . In turn, if $u \in K_c \cap (K_- \cup K_+)'$, then there exists $\{u_n\}_{n=1}^\infty \subseteq (u, \infty) \cap (K_- \cup K_+)$ such that

$u = \lim_{n \rightarrow \infty} u_n$. Hence by 2° and monotonicity of φ we must have $\varphi(s) > u$ for all $s > u$, which means that $\varphi(u+) \geq u > t = \varphi(u-)$, a contradiction.

Let us fix real $t \geq 0$. If $t = 0$, then Step 1 and (9) imply that $\varphi(0) = r(A_{[0]}) = 0$. In the rest of the proof we assume that $t > 0$.

STEP 3. If $t \in K_+$, then $\varphi(t) = r(A_{[t]}) = t$.

This can be deduced from (11), $K_+ \subseteq \{a_n : n \geq 1\}$ and 2°.

STEP 4. If $t \in K_-$, then $\varphi(t) = r(A_{[t]})$.

Using 2° and Step 2, we can show that $\varphi(s) > \varphi(t+) = t$ for all $s \in (t, \infty)$, which yields $(\varphi(t), t] \cap \varphi([0, \infty)) = \emptyset$. This and $\{a_n\}_{n=1}^\infty \subseteq \varphi([0, \infty))$ lead to $(\varphi(t), t] \cap \{a_n : n \geq 1\} = \emptyset$. Applying (11), we get $r(A_{[t]}) \leq \varphi(t)$. However $\varphi(t) \in K_r \subseteq \{a_n : n \geq 1\}$, which together with (11) and $\varphi(t) < t$ gives $r(A_{[t]}) \geq \varphi(t)$.

STEP 5. If $\alpha \in [0, t)$ and $\varphi(s) = s$ for every $s \in (\alpha, t)$, then $\varphi(t) = r(A_{[t]}) = t$.

Indeed, by $\overline{\{a_n : n \geq 1\}} = \overline{\varphi([0, \infty))}$, there exists $\{k_n\}_{n=1}^\infty \subseteq \{1, 2, \dots\}$ such that $\lim_{n \rightarrow \infty} a_{k_n} = t$ and $a_{k_n} < t$ for all integers $n \geq 1$. By virtue of (9) and (11), we have $r(A_{[t]}) = t$. In turn, Step 1 and monotonicity of φ implies $\varphi(t) = t$, as desired.

STEP 6. If $t \in K_c \cap (K_- \cup K_+)'$, then $\varphi(t) = r(A_{[t]}) = t$.

Suppose first that φ is continuous on an interval $(t - \varepsilon, t)$ for some $\varepsilon > 0$. We claim that $\varphi(t) = t$. Indeed, since there exists a sequence $\{u_n\}_{n=1}^\infty \subseteq (t - \varepsilon, t) \cap (K_- \cup K_+)$ converging to t , we deduce from 2° that $\varphi(u_n+) = u_n$ tends to t as $n \rightarrow \infty$. This and $t \in K_c$ lead to $\varphi(t) = t$. Next, in view of 4°, we get $\varphi(s) = s$ for all $s \in (t - \varepsilon, t)$, and so Step 5 gives the conclusion. In the other case there exists a strictly increasing sequence $\{t_n\}_{n=1}^\infty \subseteq K_- \cup K_+$ tending to t . Condition 2° accompanied with Steps 3 and 4 implies that

$$t_n = \varphi(t_n+) \leq \varphi(t_{n+1}) = r(A_{[t_{n+1}]}) \leq r(A_{[t]}), \quad n \geq 1.$$

Letting n tend to ∞ , and applying $t \in K_c$ and (9) proves Step 6.

STEP 7. If $t \in K_c \setminus (K_- \cup K_+)'$, then $\varphi(t) = r(A_{[t]})$.

Indeed, there exists a nonempty interval $J = (\alpha, \beta)$ containing t and such that φ is continuous on J . By virtue of 4°, we can reduce the proof of Step 7 to considering the following two cases:

- (a) $\varphi(s) = \varphi(t)$ for all $s \in J$,
- (b) $\varphi(s) = s$ for all $s \in (\alpha, t)$.

If (a) holds, then $\varphi(t) \in K_r$, which by Step 2, monotonicity of φ , 1° and 2° lead either to $(\varphi(t), \infty) \cap \varphi([0, \infty)) = \emptyset$ or to $(\varphi(t), u) \cap \varphi([0, \infty)) = \emptyset$, where $u \in [\beta, \infty) \cap (K_- \cup K_+)$. Applying (11), Step 1 and the inclusions $K_r \subseteq \{a_n : n \geq 1\} \subseteq \varphi([0, \infty))$, we get $\varphi(t) = r(A_{[t]})$. The case (b) can be deduced from Step 5. \square

In view of the proof of Theorem 10 it is clear that in general for a given transformation φ there may exist many positive selfadjoint operators A such that $\varphi(s) = r(A_{[s]})$ for every $s \in [0, \infty)$. It is also worth mentioning that none of the conditions 1°-4° can be removed from Theorem 10 without affecting its conclusion.

Arguing as in the proof of Theorem 10, we can obtain the following result.

PROPOSITION 11. *Suppose that $\varphi: [0, \infty) \rightarrow [0, \infty)$ is an arbitrary function. If φ is monotonically increasing, then φ satisfies the conditions 1°-4° of Theorem 10 if and only if the following two conditions hold:*

- (i) for every $x \in [0, \infty)$, $\varphi(x) \leq x$,
- (ii) for every $x \in [0, \infty)$, if $\varphi(x) < x$, then $\varphi(v) = \varphi(x)$ for all $v \in [\varphi(x), x]$.

If φ has left-hand and right-hand limits at each point of $[0, \infty)$ and satisfies the conditions 1°, 2° and 4° of Theorem 10, then φ is monotonically increasing.

4. Examples of discontinuity of $t \mapsto r(A_{[t]})$.

EXAMPLE 12. For normal operators A , the function $t \mapsto r(A_{[t]})$ may have either left-hand or right-hand discontinuity (but never simultaneously, cf. Proposition 8). Indeed, if α , β and γ are positive real numbers such that $\alpha < \beta < \gamma$, and $a = \{a_n\}_{n=1}^\infty$ is a sequence whose entries form a dense subset of $[0, \alpha] \cup [\beta, \gamma]$, then by (11) the function $t \mapsto r((M_a)_{[t]})$ is left-discontinuous at β whenever $a_n = \beta$ for some $n \geq 1$, and right-discontinuous at β otherwise.

EXAMPLE 13. Let \mathbb{Z} stand for the set of all integers. Denote by $\ell^2(\mathbb{Z})$ the Hilbert space of all square summable two-sided complex sequences. For $u \in [0, 1]$, we define the bounded linear operator T_u on $\ell^2(\mathbb{Z})$ by

$$T_u(\dots, x_{-1}, \boxed{x_0}, x_1, \dots) = (\dots, x_{-2}, \boxed{x_{-1}}, ux_0, ux_1, \dots), \quad \{x_j\}_{j \in \mathbb{Z}} \in \ell^2(\mathbb{Z}),$$

where the boxed entries occupy the zeroth position. We show that

$$(12) \quad r((T_u)_{[t]}) = \begin{cases} 0 & \text{if } t \in [0, u), \\ 1 & \text{if } t \in [u, \infty). \end{cases}$$

Indeed, note first that if $x \in \ell^2(\mathbb{Z})$ is nonzero, then $\|T_u x\| \geq u\|x\|$, which, when iterated, leads to $r(T_u, x) \geq u$. Hence $\mathcal{H}_{T_u}(t) = \{0\}$ for all $t \in [0, u)$. On the other hand if $t \in [u, \infty)$, then $e_n \in \mathcal{H}_{T_u}(t)$ for every $n \in \mathbb{Z}$, where $\{e_n\}_{n \in \mathbb{Z}}$ is the standard “0-1” orthonormal basis of $\ell^2(\mathbb{Z})$. This and Lemma 1 (iii) imply $r((T_u)_{[t]}) = r(T_u)$. However, $\|T_u^k\| = 1$ for all integers $k \geq 1$, which gives $r(T_u) = 1$. This justifies (12).

Take now $u, v \in \mathbb{R}$ such that $0 \leq u < v$. Since $\mathcal{H}_{\alpha A}(t) = \mathcal{H}_A(t/\alpha)$ and consequently $r((\alpha A)_{[t]}) = \alpha r(A_{[t/\alpha]})$ for $A \in \mathcal{B}(\mathcal{H})$ and $\alpha \in (0, \infty)$, we deduce from (12) that the operator $S_{u,v} \stackrel{\text{def}}{=} vT_{u/v} \in \mathcal{B}(\ell^2(\mathbb{Z}))$ has the following property

$$(13) \quad r((S_{u,v})_{[t]}) = \begin{cases} 0 & \text{if } t \in [0, u), \\ v & \text{if } t \in [u, \infty). \end{cases}$$

EXAMPLE 14. Take $u \in [0, 1)$ and a sequence $\{u_n\}_{n=1}^\infty \subseteq (u, 1)$ tending to u . Set $T = \bigoplus_{n=1}^\infty T_{u_n}$, where $T_{u_n} \in \mathcal{B}(\ell^2(\mathbb{Z}))$ are as in Example 13. The Hilbert space \mathcal{H} on which the bounded operator T acts is the orthogonal sum of \aleph_0 copies of $\ell^2(\mathbb{Z})$. We show that

$$(14) \quad r(T_{[t]}) = \begin{cases} 0 & \text{if } t \in [0, u], \\ 1 & \text{if } t \in (u, \infty). \end{cases}$$

For this, observe that if $f = \bigoplus_{n=1}^\infty f_n \in \mathcal{H}$ and $f_k \neq 0$ for some $k \geq 1$, then

$$(15) \quad r(T, f) \geq r(T_{u_k}, f_k) \geq u_k > u.$$

The first inequality can be proved similarly to (2), while the second one is justified in Example 13. Condition (15) implies that $\mathcal{H}_T(t) = \{0\}$ for all $t \in [0, u]$. Fix now $t \in (u, \infty)$ and choose an integer $k_0 \geq 1$ such that $u_{k_0} \leq t$. Denote by \mathcal{X} the set of all $f = \bigoplus_{n=1}^\infty f_n \in \mathcal{H}$ such that $f_n = 0$ for each $n \neq k_0$, and $f_{k_0} \in \{e_j : j \in \mathbb{Z}\}$, where e_j 's are as in Example 13. Since $r(T_{u_{k_0}}, e_j) = u_{k_0}$ for all integers j , we see

that $\mathcal{X} \subseteq \mathcal{H}_T(t)$. Hence, again by Lemma 1 (iii), the linear span \mathcal{E} of \mathcal{X} is contained in $\mathcal{H}_T(t)$. Consequently

$$1 = \|T\| \geq r(T) \geq r(T|_{\mathcal{E}}) = r(T_{u_{k_0}}) = 1,$$

the last equality being shown in Example 13. This completes the proof of (14).

Employing (14) and the same ‘‘scaling’’ procedure as in Example 13, we construct for each $(u, w) \in \mathbb{R}^2$ with $0 \leq u < w$ the operator $T \in \mathbf{B}(\mathcal{H})$ such that

$$(16) \quad r(T_{[t]}) = \begin{cases} 0 & \text{if } t \in [0, u], \\ w & \text{if } t \in (u, \infty). \end{cases}$$

LEMMA 15. *Let A_j be an operator with invariant domain acting in a Hilbert space \mathcal{H}_j , $j = 1, 2, \dots, n$. If $A = \bigoplus_{j=1}^n A_j$, then*

- 1° $r(A, f) = \max_{j=1, \dots, n} r(A_j, f_j)$ for all $f = \bigoplus_{j=1}^n f_j \in \mathcal{D}(A)$,
- 2° $\mathcal{H}_A(t) = \bigoplus_{j=1}^n \mathcal{H}_{A_j}(t)$ for all $t \in [0, \infty)$,
- 3° $r(A_{[t]}) = \max_{j=1, \dots, n} r((A_j)_{[t]})$ for all $t \in [0, \infty)$ such that $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$.

PROOF. Arguing as in (2) and using Lemma 1 (i) we obtain 1°. Condition 2° follows from 1°, while 3° is a consequence of 2°. \square

EXAMPLE 16. Take $u, v, w \in \mathbb{R}$ such that $0 \leq u < v < w$. Let $T \in \mathbf{B}(\mathcal{H})$ satisfy (16) and $S_{u,v}$ be as in (13). If $A = T \oplus S_{u,v}$, then by Lemma 15 we have

$$(17) \quad r(A_{[t]}) = \begin{cases} 0 & \text{if } t \in [0, u), \\ v & \text{if } t = u, \\ w & \text{if } t \in (u, \infty). \end{cases}$$

This means that the function $s \mapsto r(A_{[s]})$ is both left- and right-discontinuous at u , and $r(A_{[t]}) > t$ for all $t \in [u, w)$. In turn, the operator $A = S_{u,v}$ satisfies (17) with $0 \leq u < v = w$, while the operator $A = T \oplus uI$ (I = the identity operator on a nonzero Hilbert space) fulfills (17) with $0 \leq u = v < w$. Finally, note that by Lemma 6 (vii) there is no bounded operator A satisfying (17) with $0 < v < u$.

5. Localoid and locally normaloid operators. We now distinguish two new classes of operators. An operator A in \mathcal{H} is said to be *localoid* if $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ and $r(A_{[t]}) \leq t$ for every real $t > 0$. This definition remains unchanged if we let t range over $[0, \infty)$. Indeed, Lemma 6 leads to $r(A_{[0]}) \leq r(A_{[t]}) \leq t$ for all real $t > 0$, and so $r(A_{[0]}) = r(A_{[0+]}) = 0$. The following fact is a consequence of Lemma 6.

PROPOSITION 17. *If A is a localoid operator in \mathcal{H} , then*

- (i) $\mathcal{H}_A(t) = \mathcal{H}_A(r(A_{[t]}))$ for all $t \in [0, \infty)$,
- (ii) $r(A_{[t]}) = r(A_{[t']})$ with $t' \stackrel{\text{def}}{=} r(A_{[t]})$ for all $t \in [0, \infty)$,
- (iii) $r(A_{[t]}) = t$ whenever $t \in [0, \infty)$ and $t = r(A, f)$ for some $f \in \mathcal{D}^\infty(A)$,
- (iv) $r(A_{[t-]}) \leq r(A_{[t]}) \leq r(A_{[t+]}) \leq t$ for all $t \in [0, \infty)$,
- (v) if $t \in [0, \infty)$ and $r(A_{[t]}) < r(A_{[t+]})$, then $r(A_{[t+]}) = t$,
- (vi) $r(A_{[t]})$ equals $r(A_{[t-]})$ or $r(A_{[t+]})$ for all $t \in [0, \infty)$.

COROLLARY 18. *If A is a localoid operator in \mathcal{H} and s, t are real numbers such that $0 \leq t \leq s$, then the following conditions are equivalent*

- (i) $\mathcal{H}_A(t) = \mathcal{H}_A(s)$,
- (ii) $\mathcal{H}_A(r(A_{[t]})) = \mathcal{H}_A(r(A_{[s]}))$,

- (iii) $r(A_{[t]}) = r(A_{[s]})$,
- (iv) $r(A_{[s]}) \leq t$.

PROOF. Apply Proposition 17 (i), the localoidity of A , and (7). \square

Closed localoid operators can be characterized in a topological manner.

LEMMA 19. *Let A be a closed operator in \mathcal{H} . Then the following conditions are equivalent*

- (i) A is localoid,
- (ii) for every $t \in (0, \infty)$, $\mathcal{H}_A(t)$ is a closed subspace of \mathcal{H} ,
- (iii) for every $t \in (0, \infty)$, $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ and $r(A_{[t]}) = r(A, f_t)$ for some $f_t \in \mathcal{H}_A(t)$.

Moreover, if (i) holds, then (ii) and (iii) are also valid for $t = 0$.

PROOF. (i) \Rightarrow (ii) Suppose that $t \in (0, \infty)$ and $f \in \overline{\mathcal{H}_A(t)}$. Since $\overline{A_{[t]}} \subseteq A$ and $\overline{A_{[t]}} \in \mathbf{B}(\overline{\mathcal{H}_A(t)})$, we obtain $\overline{A_{[t]}} = A|_{\overline{\mathcal{H}_A(t)}}$ and thus $\overline{\mathcal{H}_A(t)} \subseteq \mathcal{D}^\infty(A)$. This yields $r(A, f) = r(\overline{A_{[t]}}|_{\overline{\mathcal{H}_A(t)}}, f) \leq r(\overline{A_{[t]}}) \leq t$, which means that $f \in \mathcal{H}_A(t)$. Hence $\mathcal{H}_A(t) = \overline{\mathcal{H}_A(t)}$. Since $\mathcal{H}_A(0) = \bigcap_{s>0} \mathcal{H}_A(s)$, we see that $\mathcal{H}_A(0)$ is closed.

(ii) \Rightarrow (i) By the closed graph theorem $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ for every $t \in (0, \infty)$. Owing to [8, Lemma 2] (see also [19, Theorem 1]), we have

$$r(A_{[t]}) = \max_{f \in \mathcal{H}_A(t)} r(A_{[t]}, f) = \max_{f \in \mathcal{H}_A(t)} r(A, f) \leq t, \quad t \in (0, \infty).$$

(i) \Rightarrow (iii) In view of implication (i) \Rightarrow (ii), $\mathcal{H}_A(t)$ is a closed subspace of \mathcal{H} , and so (iii) is a direct consequence of [8, Lemma 2] applied to $A_{[t]}$ ($t \in [0, \infty)$).

(iii) \Rightarrow (i) As $f_t \in \mathcal{H}_A(t)$, we get $r(A_{[t]}) = r(A, f_t) \leq t$ for all real $t > 0$. \square

An operator A in \mathcal{H} is called a *locally normaloid* operator if $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ and $\|A_{[t]}\| \leq t$ for all real $t > 0$. We may again admit $t = 0$ without harming the definition. It is clear that every locally normaloid operator is localoid. The reverse implication is no longer true (consider a nonzero quasinilpotent $A \in \mathbf{B}(\mathcal{H})$).

Locally normaloid operators can be characterized as follows.

LEMMA 20. *If A is an operator in \mathcal{H} , then the following conditions are equivalent*

- (i) A is locally normaloid,
- (ii) $\|Af\| \leq r(A, f)\|f\|$ for all $f \in \mathcal{D}^\infty(A)$,
- (iii) the sequence $\{\|(tA)^n f\|\}_{n=0}^\infty$ is monotonically decreasing for all $f \in \mathcal{H}_{tA}(1)$ and $t \in (0, \infty)$,
- (iv) $\sup_{n \geq 1} \|(tA)^n f\| \leq \|f\|$ for all $f \in \mathcal{H}_{tA}(1)$ and $t \in (0, \infty)$.

PROOF. (i) \Rightarrow (ii) Fix $f \in \mathcal{D}^\infty(A)$. Suppose that $t \stackrel{\text{def}}{=} r(A, f) < \infty$. Then $f \in \mathcal{H}_A(t)$, and consequently by (i) we have $\|Af\| = \|A_{[t]}f\| \leq t\|f\| = r(A, f)\|f\|$. The case $r(A, f) = \infty$ is obvious.

(ii) \Rightarrow (iii) Fix $f \in \mathcal{D}^\infty(A)$ and $t \in (0, \infty)$ such that $r(tA, f) \leq 1$. It follows from Lemma 1 (ii) that $r(A, A^n f) = r(A, f) \leq 1/t$ for all $n \geq 0$. Thus by (ii) we have $\|A^{n+1}f\| = \|A(A^n f)\| \leq (1/t)\|A^n f\|$ for all $n \geq 0$, which gives $\|(tA)^{n+1}f\| \leq \|(tA)^n f\|$ for all $n \geq 0$.

(iii) \Rightarrow (iv) Evident.

(iv) \Rightarrow (i) If $t \in (0, \infty)$ and $f \in \mathcal{H}_A(t)$, then $r(\frac{1}{t}A, f) \leq 1$ and hence, by (iv), $\|(1/t)Af\| \leq \|f\|$. This completes the proof. \square

We now show that under some extra assumption localoidity may imply local normaloidity. Following [1], we say that an operator A in \mathcal{H} is *restriction-normaloid* if for every linear subspace \mathcal{E} of $\mathcal{D}(A)$ such that $A|_{\mathcal{E}} \in \mathbf{B}(\mathcal{E})$, the equality $\|A|_{\mathcal{E}}\| = r(A|_{\mathcal{E}})$ holds (equivalently: $\|A|_{\mathcal{E}}\| = \sup\{|\langle Af, f \rangle| : f \in \mathcal{E}, \|f\| = 1\}$; cf. [11, §2.5.4, Theorem 2]). Note that for $A \in \mathbf{B}(\mathcal{H})$, our definition agrees with the one in [1].

THEOREM 21. *Let A be an operator in \mathcal{H} . Then the following conditions are equivalent*

- (i) A is locally normaloid,
- (ii) A is localoid and restriction-normaloid.

Moreover, if A is closed, then (i) is equivalent to

- (iii) $\mathcal{H}_A(t)$ is a closed subspace of \mathcal{H} and¹ $\|A|_{\mathcal{H}_A(t)}\| = r(A|_{\mathcal{H}_A(t)})$ for all $t \in (0, \infty)$.

PROOF. (i) \Rightarrow (ii) Evidently A is localoid. Let \mathcal{E} be a linear subspace of $\mathcal{D}(A)$ such that $A|_{\mathcal{E}} \in \mathbf{B}(\mathcal{E})$. It follows from the implication (i) \Rightarrow (ii) of Lemma 20 that

$$\|A|_{\mathcal{E}}f\| = \|Af\| \leq r(A, f)\|f\| = r(A|_{\mathcal{E}}, f)\|f\| \leq r(A|_{\mathcal{E}})\|f\|, \quad f \in \mathcal{E},$$

which leads to $\|A|_{\mathcal{E}}\| \leq r(A|_{\mathcal{E}})$. Hence $\|A|_{\mathcal{E}}\| = r(A|_{\mathcal{E}})$.

(ii) \Rightarrow (i) By our assumption we have $\|A|_{\mathcal{H}_A(t)}\| = r(A|_{\mathcal{H}_A(t)}) \leq t$ for all $t \in (0, \infty)$.

(i) \Leftrightarrow (iii) This can be deduced from Lemma 19 and implication (i) \Rightarrow (ii). \square

Note that the assumption on restriction-normaloidity can not be dropped in Theorem 21 (ii) (consider a nonzero quasinilpotent operator). It is an open question whether there exist restriction-normaloid operators which are not locally normaloid.

The following simple fact is stated without proof.

LEMMA 22. *If A is a localoid (resp. locally normaloid) operator in \mathcal{H} , then*

- (i) A^k is localoid (resp. locally normaloid) for every integer $k \geq 1$,
- (ii) $A|_{\mathcal{E}}$ is localoid (resp. locally normaloid) for every linear subspace \mathcal{E} of $\mathcal{D}(A)$ such that $A(\mathcal{E}) \subseteq \mathcal{E}$.

We show by example that both localoidity and local normaloidity are not preserved when taking adjoints (see also Example 26).

EXAMPLE 23. Let A be the bounded weighted shift on \mathcal{H} given by $Ae_n = \lambda_n e_{n+1}$ for all $n \geq 0$, where $\{\lambda_n\}_{n=0}^{\infty}$ is a sequence of positive real numbers and $\{e_n\}_{n=0}^{\infty}$ is an orthonormal basis of \mathcal{H} . We show that A^* is localoid if and only if $r(A) = 0$. The “if” part is plain due to $r(A) = r(A^*)$. For the “only if” part, note that $A^{*(n+1)}e_n = 0$ for all $n \geq 0$, which implies that $e_n \in \mathcal{H}_{A^*}(0)$ for all $n \geq 0$. By Lemma 1, $\mathcal{H}_{A^*}(0)$ is dense in \mathcal{H} and consequently by Lemma 19, $\mathcal{H}_{A^*}(0) = \mathcal{H}$. This yields $r(A) = r(A^*) = r((A^*)_{[0]}) \leq 0$, which completes the proof. Both possibilities $r(A) > 0$ and $r(A) = 0$ can occur in the context of weighted shifts (cf. [27]).

Following [34], we indicate two subclasses of locally normaloid operators.

PROPOSITION 24. *Every operator A in \mathcal{H} satisfying the inequality*

$$(18) \quad (-1)^j \Re \langle A^j f, f \rangle \geq 0, \quad f \in \mathcal{D}^{\infty}(A), \quad j = 1, 2, \dots,$$

is locally normaloid.

¹ Note that by the closed graph theorem, $A|_{\mathcal{H}_A(t)} \in \mathbf{B}(\mathcal{H}_A(t))$.

PROOF. By the Protter inequality (cf. [24, Theorem 1]), we have

$$\|Af\| \leq b_n \|f\|^{n/(n+1)} \|A^{n+1}f\|^{1/(n+1)}, \quad f \in \mathcal{D}^\infty(A), \quad n = 1, 2, \dots,$$

where $b_n^2 = (n+1)n^{-n/(n+1)}$. Letting n tend to ∞ gives $\|Af\| \leq r(A, f)\|f\|$ for all $f \in \mathcal{D}^\infty(A)$. This implies that A is locally normaloid (cf. Lemma 20). \square

Recall that an operator A in \mathcal{H} is said to be *paranormal* if $\|Ah\|^2 \leq \|A^2h\|\|h\|$ for all $h \in \mathcal{D}(A^2)$ (cf. [16, 22, 34, 11]). The following fact is a direct consequence of [34, Proposition 1] and [32, Lemma 8].

PROPOSITION 25. *If A is a paranormal operator in \mathcal{H} , then A is locally normaloid and for all $f \in \mathcal{D}^\infty(A)$, the sequence $\{\|A^n f\|^{1/n}\}_{n=1}^\infty$ is convergent in $[0, \infty]$.*

EXAMPLE 26. We give an example of a bounded locally normaloid operator, which is not paranormal and which fails to fulfill (18). Assume that \mathcal{H} is a separable infinite dimensional Hilbert space with an orthonormal basis $\{e_n\}_{n=0}^\infty$. Let $\{\varepsilon_n\}_{n=0}^\infty$ be a sequence of nonnegative real numbers which converges to 0. Then evidently

$$(19) \quad \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{j=0}^n \varepsilon_j = 0.$$

Let $A \in \mathbf{B}(\mathcal{H})$ be the weighted shift given by $Ae_n = e^{-\varepsilon_n} e_{n+1}$ for all $n \geq 0$. Since $\sup_{n \geq 0} e^{-\varepsilon_n} = 1$, we see that $\|A\| = 1$. It follows from (19) that

$$(20) \quad r(A, e_j) = \limsup_{n \rightarrow \infty} e^{-\frac{\varepsilon_j + \dots + \varepsilon_{j+n-1}}{n}} = 1, \quad j \geq 0.$$

We claim that $r(A, f) = 1$ for all $f \in \mathcal{H} \setminus \{0\}$. Indeed, the inequality “ \leq ” follows from $r(A, f) \leq r(A) \leq \|A\| = 1$. For the opposite inequality, write $f = \sum_{j=0}^\infty \alpha_j e_j$ with nonzero $\{\alpha_j\}_{j=0}^\infty \in \ell^2$. Let p be such that $\alpha_p \neq 0$. Computing as in (2) and applying (20) we get $r(A, f) \geq r(A, \alpha_p e_p) = 1$, which proves our claim. By contractivity of A this implies that $\|Af\| \leq \|f\| = r(A, f)\|f\|$ for all $f \in \mathcal{H}$. Hence, by Lemma 20, A is locally normaloid.

To see that A does not satisfy (18), take $f = \sum_{k=0}^\infty \frac{1}{2^k} e_k$ and compute

$$(-1)^j \Re \langle A^j f, f \rangle = \left(-\frac{1}{2}\right)^j \sum_{k=0}^\infty e^{-(\varepsilon_k + \dots + \varepsilon_{k+j-1})} \left(\frac{1}{2}\right)^{2k}, \quad j \geq 1.$$

To disprove paranormality of A we have to assume that the sequence $\{\varepsilon_n\}_{n=0}^\infty$ is not monotonically decreasing. Then A can not be paranormal because otherwise

$$e^{-2\varepsilon_n} = \|Ae_n\|^2 \leq \|A^2e_n\| = e^{-\varepsilon_n} e^{-\varepsilon_{n+1}}, \quad n \geq 0,$$

which is a contradiction.

6. Normality from cyclic normality. In this section we investigate the question of essential normality of an operator whose all cyclic (or multicyclic) parts are essentially normal. This question was settled by Nussbaum in the case of symmetric operators (cf. [20, Theorem 1]). We begin with an auxiliary result.

PROPOSITION 27. *Let A be a closed operator in \mathcal{H} and let \mathcal{X} be the set of all vectors $h \in \mathcal{H}$ for which there exists a closed linear subspace \mathcal{K} of \mathcal{H} reducing A to a normal operator and containing h . If \mathcal{X} is total in \mathcal{H} , then A is normal.*

PROOF. Note first that A is densely defined. One can deduce from [30, Theorem] (e.g. making use of its “moreover” part) that there exists the greatest closed linear subspace of \mathcal{H} which reduces A to a normal operator; denote it by \mathcal{H}_n . It is now clear that $\mathcal{X} \subseteq \mathcal{H}_n$, which yields $\mathcal{H}_n = \mathcal{H}$. The proof is complete. \square

Recall that $e \in \mathcal{D}^\infty(A)$ is an *analytic vector* of an operator A in \mathcal{H} if there exists a constant $r > 0$ such that $\sum_{n=0}^\infty \|A^n e\| r^n/n! < \infty$. A densely defined operator A in \mathcal{H} is said to be *subnormal* if there exists a complex Hilbert space $\mathcal{K} \supseteq \mathcal{H}$ (an isometric embedding) and a normal operator N in \mathcal{K} such that $\mathcal{D}(A) \subseteq \mathcal{D}(N)$ and $Ah = Nh$ for all $h \in \mathcal{D}(A)$. A densely defined operator A in \mathcal{H} is called *hyponormal* if $\mathcal{D}(A) \subseteq \mathcal{D}(A^*)$ and $\|A^*h\| \leq \|Ah\|$ for all $h \in \mathcal{D}(A)$.

THEOREM 28. *If A is a densely defined operator in \mathcal{H} with invariant domain, then each of the following three conditions implies the essential normality of A :*

- (a) $\mathcal{D}(A)$ consists of analytic vectors of A and for every $e \in \mathcal{D}(A)$, the operator $A|_{\mathcal{D}_e}$ is essentially normal in $\mathcal{H}_e \stackrel{\text{def}}{=} \overline{\mathcal{D}_e}$, where $\mathcal{D}_e \stackrel{\text{def}}{=} \text{lin}\{A^n e : n \geq 0\}$,
- (b) A is hyponormal and the set of all vectors $e \in \mathcal{D}(A)$ for which the operator $A|_{\mathcal{D}_e}$ is essentially normal in \mathcal{H}_e is total in \mathcal{H} ,
- (c) for every finite subset \mathcal{E} of $\mathcal{D}(A)$, the operator $A|_{\mathcal{D}_\mathcal{E}}$ is essentially normal in $\mathcal{H}_\mathcal{E} \stackrel{\text{def}}{=} \overline{\mathcal{D}_\mathcal{E}}$, where $\mathcal{D}_\mathcal{E} \stackrel{\text{def}}{=} \text{lin}\{A^n e : n \geq 0, e \in \mathcal{E}\}$.

PROOF. (a) If $e \in \mathcal{D}(A)$, then $\|A^n e\|^2 = \int_0^\infty t^n \langle E_e(dt)e, e \rangle$ for all integers $n \geq 0$, where E_e is the spectral measure of $(A|_{\mathcal{D}_e})^*(A|_{\mathcal{D}_e})^-$. By [29, Theorem 7], the operator A is subnormal. This implies that A is closable. Since, as easily checked, [30, Corollary 1] remains valid for hyponormal operators, and each subnormal operator is hyponormal, we see that \mathcal{H}_e reduces \bar{A} to a normal operator for every $e \in \mathcal{D}(A)$. This means that the operator \bar{A} satisfies all the assumptions of Proposition 27. Hence \bar{A} is normal, as desired.

(b) Argue as in the proof of (a).

(c) Since A is subnormal if and only if $A|_{\mathcal{D}_\mathcal{E}}$ is subnormal for every finite subset \mathcal{E} of $\mathcal{D}(A)$ (cf. [29, Theorem 3]), we deduce from our assumptions that A is subnormal. Applying [30, Corollary 1] and Proposition 27 as in the proof of (a) completes the proof of Theorem 28. \square

7. All this for locally algebraic and compact operators. Below, $\mathbb{K}[X]$ stands for the ring of all polynomials in indeterminate X with coefficients in a field \mathbb{K} . An operator A in \mathcal{H} is called *algebraic* if there exists a nonzero polynomial $p \in \mathbb{C}[X]$ such that $p(A)f = 0$ for all $f \in \mathcal{D}(p(A)) \stackrel{\text{def}}{=} \mathcal{D}(A^{\deg p})$. Recall that there are algebraic operators with invariant domains which are not closable (cf. [21, Example 3.2]). For the sake of self-containedness, we include the proof of the following basic fact about algebraic operators.

LEMMA 29. *Let $p_1, \dots, p_m \in \mathbb{K}[X]$ be a collection of pairwise relatively prime polynomials, \mathcal{X} be a linear space over \mathbb{K} and $A : \mathcal{X} \rightarrow \mathcal{X}$ be a linear mapping such that $p_1(A) \cdots p_m(A) = 0$. Then*

- 1° all the spaces $\mathcal{N}(p_j(A))$, $j = 1, \dots, m$, are invariant for A ,
- 2° $\mathcal{X} = \mathcal{N}(p_1(A)) \dot{+} \cdots \dot{+} \mathcal{N}(p_m(A))$, where $\dot{+}$ denotes the direct sum,
- 3° $\mathcal{N}(p_{i_1}(A) \cdots p_{i_s}(A)) = \mathcal{N}(p_{i_1}(A)) \dot{+} \cdots \dot{+} \mathcal{N}(p_{i_s}(A))$ for all finite sequences of integers $1 \leq i_1 < \cdots < i_s \leq m$,
- 4° $\mathcal{N}(p_j(A)) \neq \{0\}$ for every $j \in \{1, \dots, m\}$ such that $\prod_{k \neq j} p_k(A) \neq 0$.

PROOF. The invariance of $\mathcal{N}(p_j(A))$ under A is readily checked. The proof of 2° is by induction on m . The case $m = 1$ is clear. Suppose 2° is valid for a fixed integer $m \geq 1$. Assume that p_1, \dots, p_{m+1} are pairwise relatively prime and $p_1(A) \cdots p_{m+1}(A) = 0$. Set $q = p_1 \cdots p_m$. By relative primeness of q and p_{m+1} there exist polynomials $u, v \in \mathbb{K}[X]$ such that $uq + vp_{m+1} = 1$ (combine Corollary III.6.4, Theorem III.3.9 and Theorem III.3.11 of [12]). Then

$$(21) \quad x = q(A)u(A)x + p_{m+1}(A)v(A)x, \quad x \in \mathcal{X}.$$

As $q(A)p_{m+1}(A) = 0$, we see that

$$(22) \quad p_{m+1}(A)v(A)x \in \mathcal{N}(q(A)) \text{ and } q(A)u(A)x \in \mathcal{N}(p_{m+1}(A)) \text{ for all } x \in \mathcal{X}.$$

Define $\mathcal{Y} = \mathcal{N}(q(A))$ and $\mathcal{Z} = \mathcal{N}(p_{m+1}(A))$. Then (21) and (22) imply that

$$(23) \quad \mathcal{X} = \mathcal{Y} \dot{+} \mathcal{Z}.$$

Since \mathcal{Y} is an invariant subspace for A , $p_1(A|_{\mathcal{Y}}) \cdots p_m(A|_{\mathcal{Y}}) = 0$ and $\mathcal{N}(p_j(A|_{\mathcal{Y}})) = \mathcal{N}(p_j(A))$ for all $j = 1, \dots, m$, we infer from the induction hypothesis that $\mathcal{Y} = \mathcal{N}(p_1(A)) \dot{+} \cdots \dot{+} \mathcal{N}(p_m(A))$. By (23) this completes the induction argument.

Applying 2° to the restriction of A to $\mathcal{V} \stackrel{\text{def}}{=} \mathcal{N}(p_{i_1}(A) \cdots p_{i_s}(A))$, we get $\mathcal{V} = \mathcal{N}(p_{i_1}(A|_{\mathcal{V}})) \dot{+} \cdots \dot{+} \mathcal{N}(p_{i_s}(A|_{\mathcal{V}}))$. Since $\mathcal{N}(p_{i_k}(A|_{\mathcal{V}})) = \mathcal{N}(p_{i_k}(A))$ for all $k = 1, \dots, s$, the proof of 3° is finished.

If $\prod_{k \neq j} p_k(A) \neq 0$ and $\mathcal{N}(p_j(A)) = \{0\}$ for some $j \in \{1, \dots, m\}$, then $p_j(A)$ is injective, which contradicts $p_j(A) \prod_{k \neq j} p_k(A) = 0$. Hence 4° is proved. \square

Let A be an algebraic operator in \mathcal{H} such that $\mathcal{D}^\infty(A) \neq \{0\}$. Since $A_{[\infty]}$ is also algebraic, we find a unique monic polynomial p of minimal degree (call it *minimal*) such that $p(A_{[\infty]}) = 0$. By the fundamental theorem of algebra, $p(z) = (z - z_1)^{n_1} \cdots (z - z_m)^{n_m}$ with unique integers $n_1, \dots, n_m \geq 1$ and complex numbers z_1, \dots, z_m such that $z_j \neq z_k$ for all $j \neq k$. Owing to Lemma 29, we have

$$(24) \quad \begin{aligned} \mathcal{D}^\infty(A) &= \mathcal{N}((A_{[\infty]} - z_1)^{n_1}) \dot{+} \cdots \dot{+} \mathcal{N}((A_{[\infty]} - z_m)^{n_m}), \\ \mathcal{N}((A_{[\infty]} - z_j)^{n_j}) &\neq \{0\}, \quad j = 1, \dots, m. \end{aligned}$$

Thus every $f \in \mathcal{D}^\infty(A)$ can be uniquely decomposed as $f = f_1 + \cdots + f_m$ with $f_j \in \mathcal{N}((A_{[\infty]} - z_j)^{n_j})$. We set $\mathbf{n}_f(A) = \{j \in \{1, \dots, m\} : f_j \neq 0\}$. If no confusion can arise, we write \mathbf{n}_f instead of $\mathbf{n}_f(A)$.

PROPOSITION 30. *Let A be an algebraic operator in \mathcal{H} . Then for every nonzero vector $f \in \mathcal{D}^\infty(A)$ the sequence $\{\|A^n f\|^{1/n}\}_{n=1}^\infty$ is convergent to $\max\{|z_j| : j \in \mathbf{n}_f\}$, where z_1, \dots, z_m and \mathbf{n}_f are as above. Moreover, if $t \geq \max\{|z_j| : 1 \leq j \leq m\}$, then $\mathcal{H}_A(t) = \mathcal{D}^\infty(A)$.*

PROOF. We start with an auxiliary fact.

SUBLEMMA. *Assume that $T: \mathcal{D} \rightarrow \mathcal{D}$ is a linear mapping on an inner product space \mathcal{D} , $f \in \mathcal{D} \setminus \{0\}$, $z \in \mathbb{C} \setminus \{0\}$ and $k \geq 1$ is an integer. If $(T - z)^k f = 0$, then there exists an integer $N \geq 0$ such that the sequence $\{\frac{1}{z^{nN}} T^n f\}_{n=1}^\infty$ is convergent to an element of $\mathcal{D} \setminus \{0\}$.*

Indeed, there exists an integer $N \geq 0$ such that $(T-z)^N f \neq 0$ and $(T-z)^j f = 0$ for all $j \geq N+1$. The case $N = 0$ is easily seen to be true. If $N \geq 1$, then

$$\begin{aligned} \frac{1}{z^n n^N} T^n f &= \frac{1}{z^n n^N} ((T-z) + z)^n f = \frac{1}{n^N} \sum_{j=0}^N \binom{n}{j} \left(\frac{T-z}{z}\right)^j f \\ &= \frac{1}{n^N} \binom{n}{N} \left(\frac{T-z}{z}\right)^N f + \frac{1}{n} \sum_{j=0}^{N-1} \frac{1}{n^{N-1}} \binom{n}{j} \left(\frac{T-z}{z}\right)^j f, \quad n \geq N. \end{aligned}$$

Since $\lim_{n \rightarrow \infty} \frac{1}{n^N} \binom{n}{N} = \frac{1}{N!}$ and each sequence $\{\frac{1}{n^{N-1}} \binom{n}{j}\}_{n=N}^{\infty}$, $j = 0, \dots, N-1$, is bounded, we get $\lim_{n \rightarrow \infty} \frac{1}{z^n n^N} T^n f = \frac{1}{N!} \left(\frac{T-z}{z}\right)^N f \in \mathcal{D} \setminus \{0\}$.

We now turn to the proof of Proposition 30. There is no loss of generality in assuming that $A = A_{[\infty]}$. The spaces $\mathcal{X}_j \stackrel{\text{def}}{=} \mathcal{N}((A - z_j)^{n_j})$, $j = 1, \dots, m$, are invariant for A and, by (24), $\mathcal{D}(A) = \mathcal{X}_1 \dot{+} \dots \dot{+} \mathcal{X}_m$. Take a nonzero $f \in \mathcal{D}(A)$. Then $f = f_1 + \dots + f_m$ with the unique vectors $f_j \in \mathcal{X}_j$. Define $\mathbf{n}_f^* = \{j \in \mathbf{n}_f : |z_j| > 0\}$. We will only handle the nontrivial case of $\mathbf{n}_f^* \neq \emptyset$. Owing to Sublemma, there exist nonnegative integers $\{N_j\}_{j \in \mathbf{n}_f^*}$ such that

$$(25) \quad g_j \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{z_j^n n^{N_j}} A^n f_j \in \mathcal{X}_j \setminus \{0\}, \quad j \in \mathbf{n}_f^*.$$

Set $r = \max\{|z_j| : j \in \mathbf{n}_f^*\}$ and $N = \max\{N_j : j \in \mathbf{n}_f^*, |z_j| = r\}$. Remark that $r > 0$. We divide \mathbf{n}_f^* into two disjoint sets

$$\begin{aligned} J_1 &= \{j \in \mathbf{n}_f^* : |z_j| = r, N_j = N\}, \\ J_2 &= \{j \in \mathbf{n}_f^* : |z_j| = r, N_j < N\} \cup \{j \in \mathbf{n}_f^* : |z_j| < r\}. \end{aligned}$$

This enables us to write (with special care for $j \in \mathbf{n}_f \setminus \mathbf{n}_f^*$)

$$(26) \quad \frac{1}{r^n n^N} A^n f = \sum_{j \in J_1} \frac{1}{r^n n^N} A^n f_j + \sum_{j \in J_2} \frac{1}{r^n n^N} A^n f_j, \quad n \geq \varkappa,$$

where $\varkappa \stackrel{\text{def}}{=} \max\{n_1, \dots, n_m\}$. Note that

$$(27) \quad \frac{1}{r^n n^N} A^n f_j = \frac{1}{n^{N-N_j}} \left(\frac{z_j}{r}\right)^n \frac{1}{z_j^n n^{N_j}} A^n f_j, \quad j \in \mathbf{n}_f^*, n \geq \varkappa.$$

If $j \in J_2$, then $\lim_{n \rightarrow \infty} \frac{1}{n^{N-N_j}} \left(\frac{z_j}{r}\right)^n = 0$, which together with (25) shows that

$$(28) \quad \text{the sequence } \left\{ \sum_{j \in J_2} \frac{1}{r^n n^N} A^n f_j \right\}_{n=1}^{\infty} \text{ tends to 0 as } n \rightarrow \infty.$$

Fix an arbitrary strictly increasing sequence $\{l_n\}_{n=1}^{\infty}$ of positive integers. Passing to a subsequence if necessary, we can assume that for each $j \in J_1$ the sequence $\left\{\left(\frac{z_j}{r}\right)^{l_n}\right\}_{n=1}^{\infty}$ is convergent to a complex number α_j of absolute value 1. Hence calling upon (26), (27), (28) and (25) we see that

$$\lim_{n \rightarrow \infty} \frac{1}{r^{l_n} l_n^N} A^{l_n} f = \sum_{j \in J_1} \alpha_j g_j.$$

Since $g_j \in \mathcal{X}_j \setminus \{0\}$ and $\alpha_j \neq 0$ for all $j \in J_1$, and $\mathcal{D}(A) = \mathcal{X}_1 \dot{+} \dots \dot{+} \mathcal{X}_m$, the above limit is nonzero. This implies that $\lim_{n \rightarrow \infty} \|A^{l_n} f\|^{1/l_n} = r$. Summarizing, we have proved that every subsequence of the sequence $\{\|A^n f\|^{1/n}\}_{n=1}^{\infty}$ admits a

subsequence which is convergent to r . Hence $\{\|A^n f\|^{1/n}\}_{n=1}^\infty$ is convergent to r . This immediately implies the “moreover” part of the conclusion. \square

In the case of algebraic operators localoidity reduces to the boundedness of $A_{[\infty]}$.

PROPOSITION 31. *If A is an algebraic operator in \mathcal{H} , then the following conditions are equivalent*

- (i) A is localoid,
- (ii) $A_{[t]} \in \mathbf{B}(\mathcal{H}_A(t))$ for every $t \in (0, \infty)$,
- (iii) $A_{[\infty]}$ is bounded.

In particular, every algebraic operator $A \in \mathbf{B}(\mathcal{H})$ is localoid.

PROOF. (i) \Rightarrow (ii) Evident.

(ii) \Rightarrow (iii) Apply Proposition 30.

(iii) \Rightarrow (i) Clearly, there is no loss of generality in assuming that $A = A_{[\infty]}$. Next, since $\bar{A}: \overline{\mathcal{D}(A)} \rightarrow \overline{\mathcal{D}(A)}$ is a bounded algebraic operator, Lemma 22 enables us to reduce the proof to the case $A \in \mathbf{B}(\mathcal{H})$. Preserving the notation from the proof of Proposition 30, we define the set $\mathfrak{m}_t = \{j \in \{1, \dots, m\}: |z_j| \leq t\}$ for $t \in (0, \infty)$. It follows from Proposition 30 that

$$\mathcal{H}_A(t) = \begin{cases} \bigoplus_{j \in \mathfrak{m}_t} \mathcal{X}_j & \text{if } \mathfrak{m}_t \neq \emptyset, \\ \{0\} & \text{if } \mathfrak{m}_t = \emptyset, \end{cases} \quad t \in (0, \infty).$$

This and part 3 $^\circ$ of Lemma 29 imply that $\mathcal{H}_A(t)$ is a closed subspace of \mathcal{H} for every $t \in (0, \infty)$. Hence, by Lemma 19, A is localoid. \square

Bounded algebraic operators on \mathcal{H} may not be locally normaloid (cf. Theorem 32). However, there are unbounded closed densely defined nilpotents and idempotents with invariant domains (cf. [21]) which are evidently not localoid.

We now formulate a criterion for essential normality of algebraic operators.

THEOREM 32. *Let A be a densely defined algebraic operator in \mathcal{H} with invariant domain. Then the following conditions are equivalent*

- (i) A is closable and \bar{A} is a bounded normal operator on \mathcal{H} ,
- (ii) A is paranormal,
- (iii) A is locally normaloid.

PROOF. We need only consider the case $\mathcal{H} \neq \{0\}$.

(i) \Rightarrow (ii) This is a well known fact (cf. [34, Proposition 3]).

(ii) \Rightarrow (iii) Apply Proposition 25.

(iii) \Rightarrow (i) One way of proving this implication is to mimic the proof of [28, Proposition 6.1] and to use Lemma 20 and the discussion preceding Proposition 30. The other possibility is to reduce our case to bounded algebraic operators satisfying condition (iii) of Lemma 20, which would enable us to apply directly [28, Proposition 6.1]. Hence, in view of Lemma 20, it suffices to show that \bar{A} is a bounded algebraic locally normaloid operator. We preserve the notation from the proof of Proposition 30. By Proposition 31, A is closable and $\bar{A} \in \mathbf{B}(\mathcal{H})$. It is clear that \bar{A} is algebraic and the minimal polynomials of A and \bar{A} coincide. It follows

from Lemma 29 that

$$(29) \quad \begin{aligned} \mathcal{D}(A) &= \mathcal{N}((A - z_1)^{n_1}) \dot{+} \cdots \dot{+} \mathcal{N}((A - z_m)^{n_m}), \\ \mathcal{H} &= \mathcal{N}((\bar{A} - z_1)^{n_1}) \dot{+} \cdots \dot{+} \mathcal{N}((\bar{A} - z_m)^{n_m}), \\ \mathcal{N}((A - z_j)^{n_j}) &\subseteq \mathcal{N}((\bar{A} - z_j)^{n_j}), \quad j = 1, \dots, m. \end{aligned}$$

Fix real $t > 0$. By Proposition 31 and Lemma 19, the space $\mathcal{H}_{\bar{A}}(t)$ is closed. Hence $\overline{\mathcal{H}_A(t)} \subseteq \mathcal{H}_{\bar{A}}(t)$. We now justify the converse inclusion. Take a nonzero vector $f = \sum_{j \in \mathfrak{n}_f(\bar{A})} f_j \in \mathcal{H}_{\bar{A}}(t)$ with the unique nonzero vectors $f_j \in \mathcal{N}((\bar{A} - z_j)^{n_j})$. Let us abbreviate $\mathfrak{n}_f(\bar{A})$ to σ . We show that the space $\mathcal{U} \stackrel{\text{def}}{=} \dot{+}_{j \in \sigma} \mathcal{N}((\bar{A} - z_j)^{n_j})$ is a range of a projection $P \in \mathbf{B}(\mathcal{H})$ (a priori not orthogonal) such that

$$P(\mathcal{D}(A)) \subseteq \dot{+}_{j \in \sigma} \mathcal{N}((A - z_j)^{n_j}).$$

If $\sigma = \{1, \dots, m\}$, then $P =$ the identity operator is the only possible choice. Otherwise, the existence of such a projection follows from the equality $\mathcal{H} = \mathcal{U} \dot{+} \mathcal{V}$, where $\mathcal{V} \stackrel{\text{def}}{=} \dot{+}_{j \notin \sigma} \mathcal{N}((\bar{A} - z_j)^{n_j})$, the closedness of the spaces \mathcal{U} and \mathcal{V} (use part 3° of Lemma 29) and (29). Since $\mathcal{D}(A)$ is dense in \mathcal{H} , there exists a sequence $\{g_n\}_{n=1}^\infty \subseteq \mathcal{D}(A)$ which converges to f . Then the sequence $\{Pg_n\}_{n=1}^\infty \subseteq \dot{+}_{j \in \sigma} \mathcal{N}((A - z_j)^{n_j})$ converges to f . According to Proposition 30, $r(A, Pg_n) \leq r(\bar{A}, f)$ for all $n \geq 1$. This shows that $\mathcal{H}_{\bar{A}}(t) \subseteq \overline{\mathcal{H}_A(t)}$. Thus we have proved that $\mathcal{H}_{\bar{A}}(t) = \overline{\mathcal{H}_A(t)}$ for all real $t > 0$. As A is a bounded locally normaloid operator, we get $\|(\bar{A})_{[t]}\| = \|\bar{A}_{[t]}\| = \|A_{[t]}\| \leq t$ for all real $t > 0$, which shows that \bar{A} is locally normaloid. \square

Regarding Theorem 32 (ii), we point out that the boundedness of algebraic paranormal operators can also be deduced from [31, Theorem 1]. The essential part of the proof of Theorem 32 consists in transferring the problem from the unbounded operator case to the bounded one and then to apply [28, Proposition 6.1] (which in fact coincides with Corollary 33 below). Though the question of characterizing normality found numerous solutions in literature (e.g. [13, 1, 2, 3, 23, 25, 18]), we have not been able to come across any result directly implying the following.

COROLLARY 33. *Every algebraic restriction-normaloid operator $A \in \mathbf{B}(\mathcal{H})$ is normal.*

PROOF. By Proposition 31, the operator A is localoid. This and Theorem 21 imply that A is locally normaloid. Applying Theorem 32 completes the proof. \square

Corollary 33 is no longer true if we drop the algebraicity assumption even though the spectrum were finite. Indeed, according to [26, Theorem 2], there exists a non-normal contraction with the prescribed finite spectrum contained in the unit circle, and as such is restriction-normaloid (cf. [14, Theorem 1]). If we do not insist that the spectrum is finite, then there are obvious examples of such operators, e.g. nonunitary isometries.

An operator A in \mathcal{H} with invariant domain is said to be *locally algebraic* if for every $e \in \mathcal{D}(A)$ there exists a nonzero polynomial $p_e \in \mathbb{C}[X]$ such that $p_e(A)e = 0$. According to [17, Lemma 14] every locally algebraic bounded operator on a Banach space is automatically algebraic. This statement is no longer true for unbounded operators, e.g. every unbounded diagonal operator, when considered on “finite” vectors, is locally algebraic but not algebraic. Below we state the locally algebraic version of Theorem 32. Proposition 30 may also be adapted to this context.

THEOREM 34. *Let A be a densely defined locally algebraic operator in \mathcal{H} with invariant domain. Then the following conditions are equivalent*

- (i) A is essentially normal,
- (ii) A is paranormal,
- (iii) A is locally normaloid.

PROOF. (i) \Rightarrow (ii) \Rightarrow (iii) Mimic the appropriate parts of the proof of Theorem 32.

(iii) \Rightarrow (i) We preserve the notation introduced in Theorem 28. Take $e \in \mathcal{D}(A)$. Then $A|_{\mathcal{D}_e}$ is a densely defined algebraic operator in \mathcal{H}_e with invariant domain. Since by Lemma 22 the operator $A|_{\mathcal{D}_e}$ is locally normaloid, we infer from Theorem 32 that $A|_{\mathcal{D}_e}$ is bounded and essentially normal in \mathcal{H}_e . This implies that e is an analytic vector of A . Applying Theorem 28 (a) completes the proof. \square

The alternative proof of Theorem 34 consists in employing Theorem 28 (c).

The following result is a counterpart of Proposition 30 for compact operators. Let $A \in \mathbf{B}(\mathcal{H})$ be a compact operator. Given an isolated point $z \in \mathbb{C}$ of the spectrum $\sigma(A)$ of A , we denote by $P_z \in \mathbf{B}(\mathcal{H})$ the Riesz projection attached to $\{z\}$ (cf. [9, Chapter VII.3]). In turn, $Q_t \in \mathbf{B}(\mathcal{H})$ stands for the Riesz projection associated with the closed set $\sigma(A) \cap \{z \in \mathbb{C}: |z| \leq t\}$, where $t \in (0, \infty)$. Define

$$\mathfrak{k}_f(A) = \{z \in \mathbb{C}: z \text{ is an isolated point of } \sigma(A) \text{ and } P_z f \neq 0\}, \quad f \in \mathcal{H}.$$

Recall that (below I stands for the identity operator on \mathcal{H})

$$(30) \quad Q_t = I - \sum_{\substack{z \in \sigma(A) \\ |z| > t}} P_z, \quad t \in (0, \infty).$$

PROPOSITION 35. *Let $A \in \mathbf{B}(\mathcal{H})$ be a compact operator. Then A is localoid and for every vector $f \in \mathcal{H}$, the sequence $\{\|A^n f\|^{1/n}\}_{n=1}^\infty$ is convergent and*

$$(31) \quad \lim_{n \rightarrow \infty} \|A^n f\|^{1/n} = \begin{cases} \max\{|z|: z \in \mathfrak{k}_f(A)\} & \text{if } \mathfrak{k}_f(A) \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, the following equality holds

$$(32) \quad \mathcal{H}_A(t) = Q_t(\mathcal{H}), \quad t \in (0, \infty).$$

PROOF. Take $f \in \mathcal{H} \setminus \{0\}$. Consider three cases.

1 $^\circ$ $\mathfrak{k}_f(A) \neq \emptyset$ and $\varkappa \stackrel{\text{def}}{=} \max\{|z|: z \in \mathfrak{k}_f(A)\} > 0$. Then, by (30), $f \in Q_\varkappa(\mathcal{H})$. Since $\sigma(A|_{Q_\varkappa(\mathcal{H})}) = \{z \in \sigma(A): |z| \leq \varkappa\}$ and $r(A|_{Q_\varkappa(\mathcal{H})}) = \varkappa$, we can apply [8, Proposition 1] to the operator $A|_{Q_\varkappa(\mathcal{H})}$, which yields $\lim_{n \rightarrow \infty} \|A^n f\|^{1/n} = \varkappa$.

2 $^\circ$ $\mathfrak{k}_f(A) \neq \emptyset$ and $\varkappa = 0$. Since now 0 is an isolated point of $\sigma(A)$ and $P_0 = I - \sum_{z \in \sigma(A) \setminus \{0\}} P_z$, we get $f \in P_0(\mathcal{H})$. Hence

$$r(A, f) = r(A|_{P_0(\mathcal{H})}, f) \leq r(A|_{P_0(\mathcal{H})}) = 0.$$

3 $^\circ$ $\mathfrak{k}_f(A) = \emptyset$. Observe first that 0 is an accumulation point of $\sigma(A)$. For otherwise $\sigma(A)$ is finite and $P_z f = 0$ for all $z \in \sigma(A)$, which implies $f = 0$, a contradiction. Take $w \in \sigma(A) \setminus \{0\}$. Since by (30), $f \in Q_{|w|}(\mathcal{H})$, we get

$$r(A, f) = r(A|_{Q_{|w|}(\mathcal{H})}, f) \leq r(A|_{Q_{|w|}(\mathcal{H})}) = |w|.$$

Letting w tend to zero completes the proof of (31).

Noticing that $\mathcal{H}_A(t) = Q_t(\mathcal{H}) = \mathcal{H}$ for all positive $t \in [r(A), \infty)$, we see that (32) has to be verified only for $t \in (0, r(A))$. This can be done with the help of (31)

via (30) and the fact that the ranges of Riesz projections $\{P_z : z \in \sigma(A), |z| > t\}$ are linearly independent. Condition (32) and Lemma 19 imply that A is localoid. \square

In view of Theorem 21 and Proposition 31, within the class of bounded algebraic operators there is no difference between the notions of a restriction-normaloid operator and a locally normaloid operator. In turn, by Proposition 35, we see that the same observation remains valid for the class of compact operators. Hence, owing to [15, Theorem 1] and Proposition 25, we obtain the following corollary.

COROLLARY 36. *A compact operator $A \in \mathbf{B}(\mathcal{H})$ is normal if and only if it is locally normaloid.*

Acknowledgement. A substantial part of this paper was written while the first and the third authors visited Kyungpook National University during the fall of 2005; they wish to thank the faculty and administration for their warm hospitality.

References

- [1] S.K. Berberian, Some conditions on an operator implying normality, *Math. Ann.* **184** (1970), 188-192.
- [2] S. K. Berberian, Some conditions on an operator implying normality. II, *Proc. Amer. Math. Soc.* **26** (1970) 277-281.
- [3] S. K. Berberian, Some conditions on an operator implying normality. III, *Proc. Japan Acad.* **46** (1970), 630-632.
- [4] H. N. Bojadziev, New applications of Bernsteins inequality to the theory of operators: a local Sinclair lemma and a generalization of the Fuglede-Putnam theorem, *Complex analysis and applications '85*, Varna, 1985, 97-104, Publ. House Bulgar. Acad. Sci., Sofia, 1986.
- [5] K. N. Boyadzhiev, Sinclair type inequalities for the local spectral radius and related topics, *Israel J. Math.* **57** (1987), 272-284.
- [6] P. L. Butzer, H. Berens, *Semi-Groups of Operators and Approximation*, Springer-Verlag, Berlin, 1967.
- [7] I. Colojoară, C. Foiaş, *Theory of generalized spectral operators*, Mathematics and its Applications, Vol. 9, Gordon and Breach, Science Publishers, New York-London-Paris, 1968.
- [8] J. Daneš, On local spectral radius, *Časopis Pěst. Mat.* **112** (1987), 177-187.
- [9] N. Dunford, J. T. Schwartz, *Linear Operators. Part I*, Wiley Classics Library, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1988.
- [10] W. G. Faris, *Self-adjoint operators*, Lecture Notes in Mathematics, vol. 433, Springer-Verlag, Berlin-New York, 1975.
- [11] T. Furuta, *Invitation to linear operators*, Taylor & Francis, Ltd., London, 2001.
- [12] T. W. Hungerford, *Algebra*, Holt, Rinehart and Winston, Inc., New York-Montreal, 1974.
- [13] I. Istrăţescu, Structure theorems for some classes of operators, *Proc. Japan Acad.* **45** (1969) 586-589.
- [14] V. I. Istrăţescu, On some normaloid operators, *Rev. Roumaine Math. Pures Appl.* **14** (1969), 1289-1293.
- [15] V. Istrăţescu, I. Istrăţescu, On normaloid operators, *Math. Z.* **105** (1968) 153-156.
- [16] V. Istrăţescu, T. Saitô, T. Yoshino, On a class of operators, *Tôhoku Math. J.* **18** (1966), 410-413.
- [17] I. Kaplansky, *Infinite abelian groups*, University of Michigan Press, Ann Arbor 1954.
- [18] K. B. Laursen, Operators with finite ascent, *Pacific Journal of Mathematics* **152** (1992), 323-336.
- [19] V. Müller, Local spectral radius formula for operators in Banach spaces, *Czechoslovak Math. J.* **113** (1988), 726-729.
- [20] A. E. Nussbaum, Quasi-analytic vectors, *Ark. Mat.* **6** (1965), 179-191.
- [21] S. Ôta, Unbounded nilpotents and idempotents, *J. Math. Anal. Appl.* **132** (1988), 300-308.
- [22] S. Ôta, K. Schmüdgen, On some classes of unbounded operators, *Integr. Equat. Oper. Th.* **12** (1989), 211-226.

- [23] S. Prasanna, Weyl's theorem and thin spectra, *Proc. Indian Acad. Sci. Math. Sci.* **91** (1982), 59-63.
- [24] M. H. Protter, Inequalities for powers of unbounded operators, *Manuscripta Math.* **28** (1979), 71-79.
- [25] C. Qiu, Paranormal operators with countable spectrum are normal operators (Chinese), *J. Math. Res. Exposition* **7** (1987), 591-594.
- [26] B. Russo, Unimodular contractions in Hilbert space, *Pacific J. Math.* **26** (1968), 163-169.
- [27] A. L. Shields, Weighted shift operators and analytic function theory, *Topics in operator theory*, pp. 49-128. Math. Surveys, No. 13, Amer. Math. Soc., Providence, R.I., 1974.
- [28] J. Stochel, Characterizations of subnormal operators, *Studia Math.* **97** (3) (1991), 227-238.
- [29] J. Stochel, F. H. Szafraniec, On normal extensions of unbounded operators. II, *Acta. Sci. Math. (Szeged)* **53** (1989), 153-177.
- [30] J. Stochel, F. H. Szafraniec, The normal part of an unbounded operator, *Nederl. Akad. Wetensch. Proc. Ser. A* **92** (1989), 495-503.
- [31] J. Stochel, F. H. Szafraniec, Algebraic operators and moments on algebraic sets, *Portugaliae Mathematica* **51** (1994), 25-45.
- [32] J. Stochel, F. H. Szafraniec, C^∞ -vectors and boundedness, *Ann. Polon. Math.* **66** (1997), 223-238.
- [33] J. Stochel, F. H. Szafraniec, Domination of unbounded operators and commutativity, *J. Math. Soc. Japan* **55** (2003), 405-437.
- [34] F. H. Szafraniec, Kato-Protter type inequalities, bounded vectors and the exponential function, *Ann. Polon. Math.* **51** (1990), 303-312.
- [35] F.-H. Vasilescu, Residually decomposable operators in Banach spaces, *Tôhoku Math. J.* **21** (1969), 509-522.
- [36] P. Vrbová, On local spectral properties of operators in Banach spaces, *Czechoslovak Math. J.* **23** (1973), 483-492.
- [37] M. Zima, On the local spectral radius of positive operators, *Proc. Amer. Math. Soc.* **131** (2003), 845-850.

INSTYTUT MATEMATYKI, UNIWERSYTET JAGIELLOŃSKI, UL. REYMONTA 4, PL-30059 KRAKÓW
E-mail address: cichon@im.uj.edu.pl

DEPARTMENT OF MATHEMATICS, COLLEGE OF NATURAL SCIENCES, KYUNGPOOK NATIONAL
UNIVERSITY, TAEGU 702-701 KOREA
E-mail address: ibjung@knu.ac.kr

INSTYTUT MATEMATYKI, UNIWERSYTET JAGIELLOŃSKI, UL. REYMONTA 4, PL-30059 KRAKÓW
E-mail address: stochel@im.uj.edu.pl